

Perturbative Q^2 -power corrections to the structure function g_1 .

B.I. Ermolaev

Ioffe Physico-Technical Institute, 194021 St.Petersburg, Russia

M. Greco

Department of Physics and INFN, University Rome III, Rome, Italy

S.I. Troyan

St.Petersburg Institute of Nuclear Physics, 188300 Gatchina, Russia

We show that $(\sim 1/(Q^2)^k)$ power corrections to the spin structure function g_1 at small x and large Q^2 are generated perturbatively from the regulated infrared divergencies. When Q^2 are small, the corrections are $\sim (Q^2)^k$. We present the explicit series of such terms as well as the formulae for their resummation. These contributions are not included in the standard analysis of the experimental data. We argue that accounting for such terms can sizably change the impact of the power corrections conventionally attributed to the higher twists.

PACS numbers: 12.38.Cy

I. INTRODUCTION

The theoretical description of the Q^2 -dependence of the structure function g_1 in perturbative QCD is mostly performed for the kinematic region of large Q^2 . However, for the phenomenological analysis of the results of the COMPASS collaboration one needs the possible knowledge of g_1 at small x and small Q^2 (see e.g. Ref. [1]). Let us remind that the conventional Standard Approach (SA) based on combining the DGLAP evolution equations[2] with phenomenological inputs for the initial parton densities[3], cannot be used for description of g_1 in this region. Strictly speaking, the SA can be applied only for large values of x ($x \sim 1$) and large Q^2 : $Q^2 \gg Q_0^2$, where Q_0^2 is the starting point of the Q^2 -evolution. Indeed the small- x region lies beyond the reach of the SA because DGLAP does not include the total resummation of $\propto \ln^k 1/x$ terms. In order to describe the experimental data at small x , in the SA one has to include singular $\propto x^{-\alpha}$ terms in the expressions for the initial parton densities. Such factors act as the leading singularities (simple poles) in the Mellin transform of $g_1(x)$ and provide g_1 with the Regge asymptotics: $g_1 \sim x^{-\alpha}$ when $x \rightarrow 0$.

On the other hand, in our approach[4, 5] based on the total resummation of the leading $\ln^k 1/x$ terms in perturbative QCD, the Regge behaviour of g_1 at $x \rightarrow 0$ appears naturally, independent of the value of Q^2 . Indeed it results from the leading singularities of the anomalous dimensions and coefficient functions, which are branching points and not simple poles. The singularities of the anomalous dimensions and coefficient functions coincide. This is very important because it guarantees the independence of Q^2 of the intercepts of g_1 . Furthermore by fitting the experimental data in our approach one does not need ad hoc singular factors $\sim x^{-\alpha}$ in the initial parton densities.

The theoretical aspects of the power corrections to the DIS structure functions were considered in Refs. [6] -[8] at large x and Q^2 . The interplay between the perturbative and non-perturbative corrections in the hard kinematics was recently considered in detail in Ref [9]. In fitting experimental data in the small- x region, the discrepancies from the SA predictions are conventionally interpreted (see e.g. Refs. [10]) as the (non-perturbative) higher twist power $(\sim (1/Q^2)^k)$ contributions. However as the small- x region is beyond the reach of SA, the real size of the higher twist corrections can be erroneously overestimated. It is clear that systematic account of power corrections $\sim (1/Q^2)^k$ can be satisfactory only when using formulas that already account for the resummation of logarithmic contributions.

We have recently proposed in Ref. [11] a generalization of our previous results[4, 5] for g_1 . Although this extension goes beyond the logarithmic accuracy we keep, it looks quite natural. In addition to the standard region of large Q^2 , this generalization can describe the small- Q^2 region, though in a model-dependent way. Our suggestion, based on the analysis of the Feynman graphs for g_1 at small x , is to replace Q^2 by $Q^2 + \mu^2$ in our previous formulas, with μ being the infrared cut-off. Such a shift also leads to replacement the variable x by $x' = x + \mu^2/2pq$. The variable x' is similar to the Nachtmann variable¹.

¹ We are grateful to L.N. Lipatov for reminding this.

In the present paper we show that regulating the infrared divergencies originates also the power corrections for g_1 at small x . They differ from the well-known power corrections[6, 7, 8] related to the resummation of the Sudakov logarithms: first, they come from the ladder Feynman graphs in the Regge kinematics where virtual gluons are not always soft; then, in contrast to Refs. [6, 7, 8, 9], we never use² the parametrization $\alpha_s = \alpha(Q^2)$ and do not consider inclusions of power terms[13] into the standard expressions for α_s . We present the explicit formulas for the total resummation of these new corrections and show that at large Q^2 they are $\sim 1/(Q^2)^k$ but in the small- Q^2 region they are $\sim (Q^2)^k$. The paper is organized as follows: in Sect. 2 we explain why there should be the infrared perturbative and non-perturbative power corrections to g_1 . We also discuss in this Sect. the difference in the infrared properties of g_1 at large and small x . In Sect. 3 we remind the essence of the model for g_1 suggested in Ref. [11] and list explicit expressions for the singlet and non-singlet g_1 at the kinematic region of small x and arbitrary Q^2 . In Sect. 4 we give in more detail than in Ref. [11] theoretical arguments in favor of those expressions. The expressions for g_1 enlisted in Sect. 3 and proved in Sect. 4 account for the total resummation of leading logarithms of x and Q^2 when Q^2 are large but they are also valid for small Q^2 . They implicitly include the power Q^2 -corrections. The explicit expressions for the power corrections are extracted from those formulas in Sect. 5. We show here that the power corrections at large and small Q^2 are quite different but the lowest twist contribution to g_1 is always leading regardless of Q^2 . Finally, Sect. 6 is for discussion.

II. ORIGIN OF THE INFRARED-DEPENDENT CONTRIBUTIONS TO g_1

It is well-known that the DIS structure functions are introduced through the hadronic tensor $W_{\mu\nu}$. In particular, the spin-dependent part of $W_{\mu\nu}$ for the electron-proton DIS is parameterized by the structure functions g_1 and g_2 :

$$W_{\mu\nu}^{spin} = v\epsilon_{\mu\nu\lambda\rho} \frac{Mq_\lambda}{pq} \left[S_\rho g_1(x, Q^2) + \left(S_\rho - \frac{Sq}{pq} \right) g_2(x, Q^2) \right] \quad (1)$$

where p, M, S are the proton momentum, mass and spin respectively, q is the virtual photon momentum. The spin structure functions $g_{1,2}$ have the non-singlet, $g_{1,2}^{NS}$, and singlet, $g_{1,2}^S$, components. In general, $g_{1,2}$ (and all other DIS structure functions) acquire both perturbative and non-perturbative QCD contributions.

In the first place, there is a non-perturbative term $W_{\mu\nu}^{NPT}$ in the region of small Q^2 . For instance, there are known examples of the lattice calculations for some structure functions (see e.g. Ref. [14]), although basically $W_{\mu\nu}^{NPT}$ is poorly known.

The other part, $W_{\mu\nu}^{PT}$, of $W_{\mu\nu}$ includes both perturbative and non-perturbative QCD contributions. The standard way to calculate the structure functions is using the factorization. According to it, $W_{\mu\nu}$ is regarded as a convolution

$$W_{\mu\nu}^{PT} = W_{\mu\nu}^q \otimes \Phi_q + W_{\mu\nu}^g \otimes \Phi_g \quad (2)$$

of the partonic tensors $W_{\mu\nu}^q, W_{\mu\nu}^g$ (where q, g label the incoming parton, i.e. a quark or a gluon respectively) and probabilities $\Phi_{q,g}$ to find this parton (quark or gluon) in the incoming hadron. The probabilities $\Phi_{q,g}$ include both perturbative and non-perturbative contributions whereas purely perturbative tensors $W_{\mu\nu}^{q,g}$ describe their x - and Q^2 -evolutions. There are no known explicit expressions for $\Phi_{q,g}$. Instead, they are approximated by the initial parton densities δq and δg defined from fitting experimental data at $Q^2 \sim 1 \text{ GeV}^2$ and $x \sim 1$. The partonic tensors $W_{\mu\nu}^q, W_{\mu\nu}^g$ evolve these densities into the region where $Q^2 \gg \mu^2$ and $x \ll 1$, with μ^2 being the starting point of the Q^2 -evolution. Such a contribution to $W_{\mu\nu}$ is called the lowest twist contribution $W_{\mu\nu}^{LT}$. Besides, there are the higher twists contributions $W_{\mu\nu}^{HT}$. They can be interpreted either in terms of more involved convolution or as essentially non-perturbative objects.

In order to calculate $W_{\mu\nu}^{q,g}$, one has to regulate infrared (IR) singularities in involved Feynman graphs. Such a regulation is different for large and small x . At large x one can use DGLAP. In DGLAP this problem is solved with assuming a non-zero virtuality μ^2 for the initial partons and imposing the ordering

$$\mu^2 < k_{1\perp}^2 < k_{2\perp}^2 < \dots < k_{n\perp}^2 < Q^2 \quad (3)$$

on the transverse momenta of the ladder partons (the numeration in Eq. (3) runs from the bottom to the top of the ladders). Eq. (3) manifests that $k_{r\perp}$ acts as an infrared cut-off for integrating over k_{r+1} . However, the ordering allows to collect the contributions that are leading at large x only. In order to account for the leading (double-logarithmic)

² It is known (see Ref. [12]) that this parameterizations fails for g_1 at small x

contributions at $x \ll 1$, the upper limit of the integration in Eq. (3) should be changed for $(p+q)^2 \approx 2pq$ and the ordering should be lifted. Without the ordering, the IR singularities in the ladder rungs are not any longer regulated automatically, so an infrared cut-off should be introduced for integration over every loop momentum. It is the reason why the μ -dependence of g_1 is getting more involved at small x . Usually the cut-off is identified with μ , the starting point of the Q^2 -evolution, though it is not mandatory. Obviously, the value of μ should be large enough to justify using the Perturbative QCD:

$$k_i^2 > \mu^2 > \Lambda_{QCD}^2. \quad (4)$$

Generally speaking, there are different ways to introduce infrared cut-offs but providing IR-divergent propagators with fictitious masses is most wide-spread. Obviously, no observables should depend on a value of μ and ways of its introducing. It means that the explicit μ -dependence in $W_{\mu\nu}^{q,g}$ should be compensated by a μ -dependence of δq , δg . However, the latter are known as phenomenological expressions containing a set of numerical parameters fixed from fitting experimental data. Those parameters are supposed to be μ -dependent, though in an unexplicit way.

Before considering the IR properties of g_1 at small x , let us discuss the well-known expression for the non-singlet g_1 in the Standard Approach:

$$g_1^{NS}(x, Q^2, \mu^2) = (e_q^2/2) \int_{-\infty}^{\infty} \frac{d\omega}{2\pi i} x^{-\omega} C_q(\omega) \delta q(\omega) \exp \left[\int_{\mu^2}^{Q^2} \frac{dk_{\perp}^2}{k_{\perp}^2} \gamma_{qq}(k_{\perp}^2, \alpha_s(k_{\perp}^2)) \right] \quad (5)$$

where e_q is the quark electric charge, C_q is the coefficient function and γ_{qq} is the DGLAP non-singlet anomalous dimension. When γ_{qq} is taken in LO, the integral in Eq. (5) is known to be $(1/b) \ln[\ln(Q^2/\mu^2)/\ln(\mu^2/\Lambda_{QCD}^2)]$, with b being the first coefficient of the Gell-Mann-Low function. The expression for the singlet g_1 looks similar, though more involved. Both the coefficient functions and the anomalous dimensions in SA are known in few first orders in α_s . It means that using Eq. (5) is theoretically based for the kinematic region where x are not far from 1 and $Q^2 \gg \mu^2 \approx 1 \text{ GeV}^2$. The expression for g_1 in Eq. (5) depends on the value of μ . This dependence is supposed to disappear when g_1 is complemented by a contribution g_1^{HT} extracted from $W_{\mu\nu}^{HT}$. In practice, the treating experimental data on Polarized DIS is carried out as follows (see e.g. the recent paper Ref. [10] and refs therein): the data are compared with g_1 of Eq. (5) and the discrepancy is attributed to the higher twists impact. However, SA for g_1 is reliable at large x . At the small- x region it should be modified. In the next section we present explicit expressions for g_1^S and g_1^{NS} replacing the DGLAP expressions in the small- x region.

III. EXPRESSIONS FOR g_1 AT SMALL x AND ARBITRARY Q^2

When $x \ll 1$, the contributions $\sim \ln^k(1/x)$ are large, so they should be accounted for to all orders in the QCD coupling. The total resummation of the leading logarithms of x for g_1 was done in Refs. [4, 5] for the region of $Q^2 \gtrsim \mu^2$. We remind that, contrary to DGLAP, the expressions for g_1 in Refs. [4, 5] are valid both for large Q^2 and for $Q^2 \sim \mu^2$. Recently, in Ref. [11] we have suggested a simple prescription to generalize those results to arbitrary values of Q^2 : in the formulas of Refs. [4, 5] Q^2 should be replaced by $Q^2 + \mu^2$. This conclusion follows from the observation that the contributions of Feynman graphs to g_1 at small x depend on Q^2 through $Q^2 + \mu^2$ only. It automatically leads to the shift $x \rightarrow x + z$, with $z = \mu^2/2pq$. As the prescription is beyond the logarithmic accuracy that we kept in our previous papers, we call it a model. The theoretical grounds of this model are given in the next Sect. According to our results, the non-singlet, g_1^{NS} , component of g_1 at the small- x region is given by the following expression:

$$g_1^{NS}(x+z, Q^2 + \mu^2) = (e_q^2/2) \int_{-\infty}^{\infty} \frac{d\omega}{2\pi i} \left(\frac{1}{z+x} \right)^{\omega} C_{NS}(\omega) \delta q(\omega) \left(\frac{Q^2 + \mu^2}{\mu^2} \right)^{H_{NS}(\omega)}, \quad (6)$$

with the new coefficient functions C_{NS} ,

$$C_{NS}(\omega) = \frac{\omega}{\omega - H_{NS}(\omega)} \quad (7)$$

and the anomalous dimensions H_{NS} ,

$$H_{NS} = (1/2) \left[\omega - \sqrt{\omega^2 - B(\omega)} \right] \quad (8)$$

where

$$B(\omega) = (4\pi C_F(1 + \omega/2)A(\omega) + D(\omega))/(2\pi^2) . \quad (9)$$

$D(\omega)$ and $A(\omega)$ in Eq. (9) are expressed in terms of $\rho = \ln(1/x)$, $\eta = \ln(\mu^2/\Lambda_{QCD}^2)$, $b = (33 - 2n_f)/12\pi$ and the color factors $C_F = 4/3$, $N = 3$:

$$D(\omega) = \frac{2C_F}{b^2 N} \int_0^\infty d\rho e^{-\omega\rho} \ln\left(\frac{\rho + \eta}{\eta}\right) \left[\frac{\rho + \eta}{(\rho + \eta)^2 + \pi^2} + \frac{1}{\rho + \eta} \right] , \quad (10)$$

$$A(\omega) = \frac{1}{b} \left[\frac{\eta}{\eta^2 + \pi^2} - \int_0^\infty \frac{d\rho e^{-\omega\rho}}{(\rho + \eta)^2 + \pi^2} \right] . \quad (11)$$

H_{NS} and C_{NS} account for DL and SL contributions to all orders in α_s and, contrary to the DGLAP phenomenology, δq does not contain singular factors. The infrared cut-off μ obeys Eq. (4). Expression (6) is valid for large Q^2 , i.e. for $Q^2 \gg \mu^2$ where $x \gg z$ and for small Q^2 , $Q^2 \leq \mu^2$ where $x \leq z$. The expression for the singlet component, g_1^S , of g_1 is more involved:

$$g_1^S = g_1^{(+)} + g_1^{(-)} , \quad (12)$$

with

$$g_1^{(\pm)} = \frac{\langle e_q^2 \rangle}{2} \int_{-\infty}^{\infty} \frac{d\omega}{2\pi i} \left(\frac{1}{z + x} \right)^\omega \left(C_q^{(\pm)} \delta q - \frac{A'}{2\pi\omega^2} C_g^{(\pm)} \delta g \right) \left(\frac{Q^2 + \mu^2}{\mu^2} \right)^{\Omega_{(\pm)}} \quad (13)$$

where $\langle e_q^2 \rangle$ stands for the sum of electric charges: $\langle e_q^2 \rangle = 10/9$ for $n_f = 4$, δq and δg are the initial quark and gluon densities.

The exponents $\Omega_{(\pm)}$ and coefficient functions $C_{q,g}^{(\pm)}$ are:

$$\Omega_{(\pm)} = \frac{1}{2} [H_{qq} + H_{gg} \pm R] . \quad (14)$$

$$\begin{aligned} C_q^{(+)} &= \frac{\omega}{RT} \left[(H_{qq} - \Omega_{(-)}) (\omega - H_{gg}) + H_{qg} H_{gq} + H_{gq} (\omega - \Omega_{(-)}) \right] , \\ C_q^{(-)} &= \frac{\omega}{RT} \left[(\Omega_{(+)} - H_{qq}) (\omega - H_{gg}) - H_{qg} H_{gq} + H_{gq} (\Omega_{(+)} - \omega) \right] , \\ C_g^{(+)} &= \frac{\omega}{RT} \left[(H_{gg} - \Omega_{(-)}) (\omega - H_{qq}) + H_{qg} H_{gq} + H_{qg} (\omega - \Omega_{(-)}) \right] \left(-\frac{A'}{2\pi\omega^2} \right) , \\ C_g^{(-)} &= \frac{\omega}{RT} \left[(\Omega_{(+)} - H_{gg}) (\omega - H_{qq}) - H_{qg} H_{gq} + H_{qg} (\Omega_{(+)} - \omega) \right] \left(-\frac{A'}{2\pi\omega^2} \right) . \end{aligned} \quad (15)$$

Here

$$R = \sqrt{(H_{qq} - H_{gg})^2 + 4H_{qg}H_{gq}} , \quad T = \omega^2 - \omega(H_{gg} + H_{qq}) + (H_{gg}H_{qq} - H_{qg}H_{gq}) \quad (16)$$

and

$$\begin{aligned} H_{qq} &= \frac{1}{2} \left[\omega + Z + \frac{b_{qq} - b_{gg}}{Z} \right] , & H_{qg} &= \frac{b_{qg}}{Z} , \\ H_{gg} &= \frac{1}{2} \left[\omega + Z - \frac{b_{qq} - b_{gg}}{Z} \right] , & H_{gq} &= \frac{b_{gq}}{Z} \end{aligned} \quad (17)$$

where

$$Z = \frac{1}{\sqrt{2}} \sqrt{(\omega^2 - 2(b_{qq} + b_{gg})) + \sqrt{(\omega^2 - 2(b_{qq} + b_{gg}))^2 - 4(b_{qq} - b_{gg})^2 - 16b_{qg}b_{gq}}} \quad (18)$$

with

$$b_{ik} = a_{ik} + V_{ik} . \quad (19)$$

The Born contributions a_{ik} are defined as follows:

$$a_{qq} = \frac{A(\omega)C_F}{2\pi} , \quad a_{gq} = -\frac{n_f A'(\omega)}{2\pi} , \quad a_{qg} = \frac{A'(\omega)C_F}{\pi} , \quad a_{gg} = \frac{4NA(\omega)}{2\pi} . \quad (20)$$

At last, non-ladder contributions are:

$$V_{ik} = \frac{m_{ik}}{\pi^2} D(\omega) , \quad (21)$$

with

$$m_{qq} = \frac{C_F}{2N} , \quad m_{gg} = -2N^2 , \quad m_{gq} = n_f \frac{N}{2} , \quad m_{qg} = -NC_F . \quad (22)$$

The additional factor $\left(-\frac{A'}{2\pi\omega^2}\right)$ in the coefficients $C_g^{(\pm)}$ in Eqs.(15), with

$$A'(\omega) = \frac{1}{b} \left[\frac{1}{\eta} - \int_0^\infty \rho \frac{d\rho e^{-\omega\rho}}{(\rho + \eta)^2} \right] , \quad (23)$$

is the small- ω estimate for the quark box diagram that dominates in the Born term relating initial gluons to the electromagnetic current. $A'(\omega)$ is the Mellin representation of the QCD running coupling α_s involved in the quark box.

Besides resummation of the leading logarithms, Eqs. (6,12) differ from DGLAP in the parametrization of α_s : the DGLAP -prescription is $\alpha_s = \alpha_s(Q^2)$ whereas in our approach α_s is replaced by A and A' defined in Eqs. (11,23). Such a difference results into a drastic difference in the form of the Q^2 -dependence of g_1 between our approach and SA: instead of the factor $(Q^2 + \mu^2)/\mu^2$ in Eqs. (6,12), the SA leads to $\ln(Q^2/\Lambda_{QCD}^2)$.

IV. THEORETICAL GROUNDS FOR THE SHIFT $Q^2 \rightarrow Q^2 + \mu^2$ IN EQS. (6) AND (12)

Both the singlet and non-singlet components of g_1 obey the following Bethe-Salpeter equation depicted in Fig. 1:

$$g_1 = g_1^{Born} + \imath \int \frac{d^4 k}{(2\pi)^4} (-2\pi\imath) \delta((q+k)^2 - m_q^2) \frac{2k_\perp^2}{(k^2 - m_q^2)^2} M(p, k) \quad (24)$$

where the δ -function (together with the factor $-2\pi\imath$) corresponds to the cut propagator of the uppermost quark with momentum k and mass m_q coupled to the virtual photon lines and $2k_\perp^2$ appear after simplifying the spin structure of the equation. $M(p, k)$ stands for the cut parton ladders and the initial parton densities. In other words, $M(p, k)$ incorporates both the initial parton densities and radiative corrections to them. This object can be called the polarized parton distribution function. In the present paper we will address it simply as the (parton) distribution, skipping other words. For the sake of simplicity we dropped unessential numerical factors ($e_q^2/2$ for f^{NS} and $< e_q^2 > /2$ for the singlet) in Eq. (24). Obviously, M in Eq. (24) cannot depend on Q^2 .

A. Prescription for the IR regularization of M

The IR-divergent contributions to M should be regulated. We follow the standard prescription and assign a fictitious mass μ to the gluons in the IR-divergent propagators. In particular, in the ladder graphs such propagators are the vertical ones. We assume that the value of μ satisfies Eq. (4). In contrast to the gluon ladders, quark ladders are IR-stable because the quark mass m_q acts as an IR cut-off. In order to use the same cut-off μ for regulating both gluon and quark IR divergences, we assume that, in addition to Eq. (4), $\mu \gg m_q$. After that m_q can be dropped. Therefore in order to regulate IR singularities, we should insert μ^2 in the IR -divergent, strut (vertical) propagators, for both quarks and gluons. The horizontal propagators (rungs) are IR-stable. This converts Eq. (24) into

$$g_1 = g_1^{Born} + \imath \int \frac{d^4 k}{(2\pi)^4} (-2\pi\imath) \delta((q+k)^2) \frac{2k_\perp^2}{(k^2 - \mu^2)^2} M(pk, k^2 + \mu^2) \quad (25)$$

B. Solving the Bethe-Salpeter equation (25)

As the kinematics $x \ll 1$ is of the Regge type, we need to express M in the Regge kinematics as well. Let us notice³ that the expressions for M can be obtained from our results for g_1 in Ref. [4, 5] with replacing the external photon virtuality $q^2 = -Q^2$ by the external quark virtuality k^2 and x by $-k^2/w\alpha$. In the first place, we focus on applying Eq. (24) to g_1^{NS} , the non-singlet part of g_1 and denote M^{NS} the involved quark distribution. The expression for M^{NS} accounting for the total resummation of leading logarithmic contributions can also be borrowed from our formula for g_1^{NS} obtained in Ref. [5]. This expression reads:

$$M^{NS}(p, k) = \int_{-\infty}^{\infty} \frac{d\omega}{2\pi i} \left(\frac{2pk}{-k^2 + \mu^2} \right)^\omega \omega f(\omega) \delta q(\omega) \left(\frac{-k^2 + \mu^2}{\mu^2} \right)^{H_{NS}(\omega)} \quad (26)$$

where $f = 8\pi^2 H_{NS}$, $\delta q(\omega)$ is the initial quark density in the ω -space and H_{NS} is given by Eq. (8). H_{NS} and f include the total resummation of leading logarithmic contributions. Similarly, the expression for the singlet distribution M^S can be obtained from our results for g_1^S in Ref. [4]. It is convenient to rewrite Eq. (25) in terms of the Sudakov variables for momentum k :

$$k = -\alpha q' + \beta p + k_\perp, \quad (27)$$

with $q' = q + xp$, so that $q'^2 \approx p^2 \approx 0$. Substituting Eq. (26) into Eq. (25), using the Sudakov variables and changing the order of the integrations, we obtain the following equation for g_1^{NS} :

$$g_1^{NS} = g_1^{Born} + \frac{1}{8\pi^2} \int_{-\infty}^{\infty} \frac{d\omega}{2\pi i} \omega f(\omega) \delta q(\omega) \int d\alpha d\beta dk_\perp^2 \frac{k_\perp^2}{(w\alpha\beta + k_\perp^2 + \mu^2)^2} \delta(w\beta + wx\alpha - w\alpha\beta - k_\perp^2 - Q^2) \left(\frac{w\alpha}{w\alpha\beta + k_\perp^2 + \mu^2} \right)^\omega \left(\frac{w\alpha\beta + k_\perp^2 + \mu^2}{\mu^2} \right)^{H_{NS}} \quad (28)$$

where we have denoted $w = 2pq$. As we consider $x \ll 1$, we can neglect $x\alpha$ compared to β . Using the δ -function for integration over β , we arrive at

$$g_1^{NS} = g_1^{Born} + \frac{1}{8\pi^2} \int_{-\infty}^{\infty} \frac{d\omega}{2\pi i} \omega f(\omega) \delta q(\omega) \int \frac{d\alpha dk_\perp^2}{\alpha Q^2 + k_\perp^2 + \mu^2} \left(\frac{w\alpha}{\alpha Q^2 + k_\perp^2 + \mu^2} \right)^\omega \left(\frac{\alpha Q^2 + k_\perp^2 + \mu^2}{\mu^2} \right)^{H_{NS}}. \quad (29)$$

The integration region in Eq. (25) is shown in Fig. 2. It is outlined by the following restrictions:

$$w \gg k_\perp^2 > 0, \quad w \gg w\alpha \gg \alpha Q^2 + k_\perp^2 + \mu^2. \quad (30)$$

Integrating over α and $k_\perp^2$ yields different contributions, depending on the ratio between αQ^2 and $k_\perp^2$. The most important contribution comes from the region D in Fig. 2. After integration over α in this region we get

$$\begin{aligned} g_1^{NS} &= g_1^{Born} + \frac{1}{8\pi^2} \int_{-\infty}^{\infty} \frac{d\omega}{2\pi i} \omega f(\omega) \delta q(\omega) \frac{1}{w} \int_{Q^2}^w \frac{dk_\perp^2}{k_\perp^2 + \mu^2} \left(\frac{w}{k_\perp^2 + \mu^2} \right)^\omega \left(\frac{k_\perp^2 + \mu^2}{\mu^2} \right)^{H_{NS}} \\ &= g_1^{Born} + \frac{1}{8\pi^2} \int_{-\infty}^{\infty} \frac{d\omega}{2\pi i} f(\omega) \delta q(\omega) \int_{Q^2 + \mu^2}^{w + \mu^2} \frac{dt}{t} \left(\frac{w}{t} \right)^\omega \left(\frac{t}{\mu^2} \right)^{H_{NS}}. \end{aligned} \quad (31)$$

The leading contribution in Eq. (31) comes from the lowest limit $t = Q^2 + \mu^2$ and gives

$$g_1^{NS} = g_1^{Born} + \frac{1}{8\pi^2} \int_{-\infty}^{\infty} \frac{d\omega}{2\pi i} \frac{f(\omega) \delta q(\omega)}{(\omega - H_{NS})} \left(\frac{w}{Q^2 + \mu^2} \right)^\omega \left(\frac{Q^2 + \mu^2}{\mu^2} \right)^{H_{NS}} \quad (32)$$

which proves the validity of the shift $Q^2 \rightarrow Q^2 + \mu^2$ suggested in Ref. [11]. Let us brush up Eq. (32). Replacing $f(\omega)$ by $8\pi^2 H_{NS}$ we see that in Eq. (32)

$$\frac{H_{NS}}{\omega - H_{NS}} = -1 + \frac{\omega}{\omega - H_{NS}}. \quad (33)$$

The first term in Eq. (33) cancels the Born contribution g_1^{Born} and the second term is the non-singlet coefficient function (see Eq. (8)). Therefore, we arrive at Eq. (6) for the non-singlet g_1^{NS} at small x and arbitrary Q^2 . Eq. (12) for the singlet g_1 in the same kinematic can be proved similarly.

³ We will consider such distributions in more detail in next our paper.

V. INFRARED POWER CORRECTIONS AT SMALL x

A. Power corrections at large Q^2

In the kinematics where $Q^2 > \mu^2$ and therefore $x > z$, the terms with $Q^2 + \mu^2$ in Eqs. (6,12) can be expanded into series in μ^2/Q^2 :

$$\begin{aligned} \left(\frac{1}{x+z}\right)^\omega \left(\frac{Q^2 + \mu^2}{\mu^2}\right)^{H_{NS}} &= \left(\frac{1}{x}\right)^\omega \left(\frac{Q^2}{\mu^2}\right)^{H_{NS}} \left[1 + \sum_{k=1} T_k^{NS}(\omega) \left(\frac{\mu^2}{Q^2}\right)^k\right], \\ \left(\frac{1}{x+z}\right)^\omega \left(\frac{Q^2 + \mu^2}{\mu^2}\right)^{\Omega_\pm} &= \left(\frac{1}{x}\right)^\omega \left(\frac{Q^2}{\mu^2}\right)^{\Omega_\pm} \left[1 + \sum_{k=1} T_k^{(\pm)}(\omega) \left(\frac{\mu^2}{Q^2}\right)^k\right] \end{aligned} \quad (34)$$

where

$$\begin{aligned} T_k^{NS} &= \frac{(-\omega + H_{NS})(-\omega + H_{NS} - 1) \dots (-\omega + H_{NS} - k + 1)}{k!}, \\ T_k^{(\pm)} &= \frac{(-\omega + \Omega_\pm)(-\omega + \Omega_\pm - 1) \dots (-\omega + \Omega_\pm - k + 1)}{k!}. \end{aligned} \quad (35)$$

It allows to rewrite Eqs. (6,12) as follows:

$$\begin{aligned} g_1^{NS}(x+z, Q^2) &= \tilde{g}_1^{NS}(x, Q^2) + \tilde{g}_1^{NS}(x/y, Q^2) \otimes \sum_{k=1} (\mu^2/Q^2)^k E_k^{NS}(y), \\ g_1^S(x+z, Q^2) &= \tilde{g}_1^S(x, Q^2) + \sum_{k=1} (\mu^2/Q^2)^k \left[\tilde{g}_1^{(+)}(x/y, Q^2) \otimes E_k^{(+)}(y) + \tilde{g}_1^{(-)}(x/y, Q^2) \otimes E_k^{(-)}(y) \right] \end{aligned} \quad (36)$$

where, using the conventional terms, $\tilde{g}_1^{NS}$ and $\tilde{g}_1^S$ can be named the non-singlet and singlet components of the lowest twist contribution to g_1 :

$$\begin{aligned} \tilde{g}_1^{NS} &= (e_q^2/2) \int_{-\infty}^{\infty} \frac{d\omega}{2\pi i} \left(\frac{1}{x}\right)^\omega C_{NS}(\omega) \delta q(\omega) \left(\frac{Q^2}{\mu^2}\right)^{H_{NS}(\omega)}, \\ \tilde{g}_1^S &= \tilde{g}_1^{(+)} + \tilde{g}_1^{(-)} = \frac{\langle e_q^2 \rangle}{2} \int_{-\infty}^{\infty} \frac{d\omega}{2\pi i} \left(\frac{1}{x}\right)^\omega \left[\left(C_q^{(+)} \left(\frac{Q^2}{\mu^2}\right)^{\Omega_{(+)}} + C_q^{(-)} \left(\frac{Q^2}{\mu^2}\right)^{\Omega_{(-)}}\right) \delta q - \right. \\ &\quad \left. \frac{A'}{2\pi\omega^2} \left(C_g^{(+)} \left(\frac{Q^2}{\mu^2}\right)^{\Omega_{(+)}} + C_g^{(-)} \left(\frac{Q^2}{\mu^2}\right)^{\Omega_{(-)}}\right) \delta g \right] \end{aligned} \quad (37)$$

and

$$E_k^{NS}(x) = \int_{-\infty}^{\infty} \frac{d\omega}{2\pi i} \left(\frac{1}{x}\right)^\omega T_k^{NS}(\omega), \quad E_k^\pm(x) = \int_{-\infty}^{\infty} \frac{d\omega}{2\pi i} \left(\frac{1}{x}\right)^\omega T_k^\pm(\omega). \quad (38)$$

The right-hand sides in Eq. (36) are the products of the power corrections and $\tilde{g}_1$. We call these corrections the infrared power corrections. The functions $\tilde{g}_1^S$, $\tilde{g}_1^{NS}$ in Eq. (37) were obtained in Refs. [4, 5] and they correspond to the lowest twist contribution. They differ from the lowest twist DGLAP expressions for g_1 by the total resummation of the leading logarithms of x and by the new parametrization of α_s given by Eqss. (11,23). They include the most important at small x contributions of the LO and NLO DGLAP formulas. When the lowest twist expressions of Eq. (37) are used for analysis of experimental data of the polarized DIS, the power series in the rhs of Eq. (36) look as new independent contributions. However, the left-hand sides of Eq. (36) account for the total resummation of these corrections. Finally, let us notice that the infrared power corrections in Eq. (36) have nothing to do with the standard parametrization $\alpha_s = \alpha_s(Q^2)$ as we do not use it.

B. Power corrections at small Q^2

When $Q^2 < \mu^2$, g_1^{NS} and g_1^S cannot be expanded similarly to Eq. (36). The power corrections for small Q^2 are different. Indeed, in this case

$$\begin{aligned} \left(\frac{1}{x+z}\right)^\omega \left(\frac{Q^2 + \mu^2}{\mu^2}\right)^{H_{NS}} &= \left(\frac{1}{z}\right)^\omega \left[1 + \sum_{k=1} T_k^{NS}(\omega) \left(\frac{Q^2}{\mu^2}\right)^k\right], \\ \left(\frac{1}{x+z}\right)^\omega \left(\frac{Q^2 + \mu^2}{\mu^2}\right)^{\Omega_\pm} &= \left(\frac{1}{z}\right)^\omega \left[1 + \sum_{k=1} T_k^{(\pm)}(\omega) \left(\frac{Q^2}{\mu^2}\right)^k\right]. \end{aligned} \quad (39)$$

It leads to the following expressions for g_1 at small Q^2 :

$$\begin{aligned} g_1^{NS}(x+z, Q^2) &= \tilde{g}_1^{NS}(z, \mu^2) + \tilde{g}_1^{NS}(z/y, \mu^2) \otimes \sum_{k=1} (Q^2/\mu^2)^k E_k^{NS}(y), \\ g_1^S(x+z, Q^2) &= \tilde{g}_1^S(z, \mu^2) + \sum_{k=1} (Q^2/\mu^2)^k \left[\tilde{g}_1^{(+)}(z/y, \mu^2) \otimes E_k^{(+)}(y) + \tilde{g}_1^{(-)}(z/y, \mu^2) \otimes E_k^{(-)}(y) \right] \end{aligned} \quad (40)$$

where the lowest twist contributions $\tilde{g}_1^{NS}$, $\tilde{g}_1^S$ are given by Eq. (37) and E_k^{NS} , E_k^S are defined in Eq. (38). Both $\tilde{g}_1^{NS}$ and $\tilde{g}_1^S$ do not depend on x and Q^2 . Instead, they depend on the total energy $(p+q)^2$ of the process and are constants when the $2pq$ is fixed. Both the x and Q^2 -dependence are now associated with the power corrections. Eqs. (36,40) show that the infrared Q^2 -corrections are different for large and small Q^2 , so that the series of Eq. (40) cannot be extrapolated into the region of small Q^2 and similarly the series in Eq. (36) cannot be extrapolated into the large Q^2 -region. Besides the terms with $(Q^2)^k$ given by Eq. (40), similar contributions can come from other sources which are beyond our control. However, the $(Q^2)^k$ -terms in Eq. (40) are multiplied by the functions $\tilde{g}_1^{NS}$ and $\tilde{g}_1^S$ which include the total resummation of the leading logarithms and therefore they are supposed to dominate, at small x , over the coefficients at the other $(Q^2)^k$ -terms.

Although the large- Q^2 expansion (34) and the small- Q^2 expansion (39) look quite similar, the power corrections to g_1^{NS} are actually different for large and small Q^2 . It is easy to check that the linear in μ^2/Q^2 term is present in the large- Q^2 expansion of Eq. (36) for g_1^{NS} while the term with Q^2/μ^2 is absent in the g_1^{NS} -expansion of Eq. (40).

VI. DISCUSSION

At the region of small x , the DGLAP ordering (3) should be lifted for accounting for leading logarithms of $1/x$, so the infrared cut-off μ should be introduced explicitly to regulate the infrared divergencies in every rung of the Feynman graphs contributing to g_1 . Similarly to DGLAP, μ can also play the role of the starting point of the Q^2 -evolution, though not obligatory. With both the perturbative and non-perturbative contributions accounted for, g_1 does not depend on μ . However, this dependence does exist when the non-perturbative contributions are neglected or accounted for implicitly through the fits for the initial parton densities. In this case the structure function g_1 depends on the value of μ and the way it has been introduced. Introducing μ as the fictitious mass inserted into the IR-divergent propagators, leads to Eqs. (6,12) suggested in Ref. [11] for g_1 . The expressions of Eqs. (6,12) include the total resummation of double-logarithms and the most important part of single-logarithms of x . They are obtained from our previous results with the shift $Q^2 \rightarrow Q^2 + \mu^2$. The theoretical grounds for such a shift are given by Eqs. (9-32). Eqs. (6,12), in contrast to DGLAP, can be used both for large and small Q^2 . Having been expanded into the series in $1/(Q^2)^k$ at large Q^2 (or into series in Q^2 at small Q^2), Eqs. (6,12) yield the power Q^2 -corrections. The series of the corrections are represented by expressions (36,40). The power series of Eqs. (36,40) for large and small Q^2 are derived from the same formulas. However after the expansion has been made, they cannot be related to each other with simply varying Q^2 . We suggest that accounting for the new source of the power contributions given by Eqs. (6,12) can sizably change the conventional analysis of the higher twists contributions to the Polarized DIS because such contributions appear in the present analysis of experimental data as a discrepancy between the experimental data and the Standard Approach predictions. Let us remind that in Ref. [15] we showed that the singular ($\sim x^{-\alpha}$) factors in the standard fits mimic the total resummations of $\ln^k x$, i.e. they have a purely perturbative origin contrary to previous common expectations. Similarly, a good portion of the commonly believed non-perturbative power corrections in the conventional analysis of experimental data can actually be of the perturbative infrared origin. However being misinterpreted as non-perturbative terms, they can mimic the power expansion in Eq. (36). In particular, Eq. (6) predicts that the power $\sim 1/(Q^2)^k$ -corrections to g_1^{NS} should appear at $Q^2 \gtrsim 1 \text{ GeV}^2$ and cannot appear at smaller values of Q^2 . It agrees with the phenomenological observations obtained in Refs. [10] from conventional analysis of



experimental data. On the other hand, Eq. (12) predicts that the similar power corrections to the singlet g_1 should be seen at greater values of Q^2 . Clearly, the use of Eqs. (6,12) for the lower twist contributions to g_1 , instead of DGLAP, would allow one to revise the impact of the genuine higher twists contributions which are known to be of the non-perturbative origin.

Finally, we would like to remind that our results explicitly depend on the infrared cut-off μ . As pointed out in Sect. II, such a dependence would vanish if analytic expressions for the probabilities $\Phi_{q,g}$ were obtained and used in Eqs. (6,12) instead of δq and δg .

VII. ACKNOWLEDGEMENTS

We are grateful to R. Windmolders and A. Korzenev for drawing our attention to the problem of the power corrections to g_1 at small x . We are also indebted to G.P. Korchemsky and S.I. Alekhin for discussing the power corrections to g_1 in the framework of the Standard Approach. The work is supported in part by the Russian State Grant for Scientific School RSGSS-5788.2006.2.

-
- [1] COMPASS coll., E.S. Ageev et al., Phys.Lett.B 612(2005)154-164; hep-ex/0701014.
 - [2] G. Altarelli and G. Parisi, Nucl. Phys.B126 (1977) 297; V.N. Gribov and L.N. Lipatov, Sov. J. Nucl. Phys. 15 (1972) 438; L.N. Lipatov, Sov. J. Nucl. Phys. 20 (1972) 95; Yu.L. Dokshitzer, Sov. Phys. JETP 46 (1977) 641.
 - [3] G. Altarelli, R.D. Ball, S. Forte and G. Ridolfi, Nucl. Phys. B496 (1997) 337; Acta Phys. Polon. B29(1998)1145; E. Leader, A.V. Sidorov and D.B. Stamenov. Phys. Rev. D73 (2006) 034023; J. Blumlein, H. Botcher. Nucl. Phys. B636 (2002) 225; M. Hirai et al. Phys. Rev. D69 (2004) 054021.
 - [4] B.I. Ermolaev, M. Greco, S.I. Troyan. Phys.Lett.B 579 (2004) 321.
 - [5] B.I. Ermolaev, M. Greco, S.I. Troyan. Nucl. Phys.B 571 (2000) 137; Nucl. Phys.B 594 (2001) 71.
 - [6] E. Stein, M. Maul, L. Mankiewicz, A. Schafer. Nucl. Phys.B 536 (1998) 318; M. Beneke, V.M. Braun. hep-ph/0010208; G.P. Korchemsky, G. Oderda, G. Sterman. hep-ph/9708346; E. Gardi, R.G. Roberts. Nucl. Phys.B 653 (2003) 227.
 - [7] G.P. Korchemsky and G. Sterman. Nucl. Phys. B 437 (1995) 415.
 - [8] G. Sterman. hep-ph/0606032.
 - [9] A.I. Karanikas, N.G. Stefanis. hep-ph/0101031; N.G. Stefanis. Phys.Lett.B504 (2001)225, Erratum-ibid.B636 (2006) 330; A.I. Karanikas, C.N. Ktorides, N.G. Stefanis. hep-ph/0201278.
 - [10] E. Leader, A.V. Sidorov, D.B. Stamenov. hep-ph/0509183; E. Leader, A.V. Sidorov, D.B. Stamenov. Phys. Rev.D 67 (2003) 074017.
 - [11] B.I. Ermolaev, M. Greco, S.I. Troyan. hep-ph/0605133.
 - [12] B.I. Ermolaev, M. Greco, S.I. Troyan. Phys.Lett.B522 (2001) 57.
 - [13] D.V. Shirkov and I.L. Solovsov. Pys. Lett. B 442 (1998) 344.
 - [14] G. Martinelli, G.C. Rossi, C.T. Sachrajda, S.R. Sharpe, M. Talevi, M. Testa. Nucl.Phys.B611 (2001)311.
 - [15] B.I. Ermolaev, M. Greco, S.I. Troyan. Phys.Lett.B 622 (2005) 93; hep-ph/0511343.

FIG. 1: The Bethe-Salpeter equation for g_1 .FIG. 2: The integration region over α and $k_{\perp}^2$ in Eq. (29).